

Eye drops

Continue with your usual anti-glaucoma drops (if any).

What to do if you experience any pain or discomfort

If you develop pain in the treated eye, take a painkiller such as paracetamol and apply a cold compress over the eye. To make a cold compress, put ice or crushed ice into a plastic bag and wrap with a moist flannel or tea towel (not dry). If it is too cold leave it for five minutes before applying it to your eye so that you don't give yourself frostbite. Apply to the closed eye, as required, for up to 15 minutes at a time.

If the pain persists, please do not hesitate to contact us.

If you have any questions or concerns please contact:

**Eye emergencies only 9am – 5pm
01342 306782**

Out of hours, bank holidays and weekends please call Ross Tilley Ward on 01342 414451

Follow-up appointment

Please make an appointment to be seen in the Glaucoma Clinic in the time period instructed.

Instructions for patients
following
Selective Laser
Trabeculoplasty (SLT)

Corneo-Plastic Unit

Please ask if you
would like this leaflet
in larger print or a
different format.

Mr Gokulan Ratnarajan - Consultant Ophthalmic Surgeon
Issue 2 - ref 0529
Approved by the Patient Information Group
Print June 2019 - Review June 2022

© Copyright QVH NHS Foundation Trust
qvh.nhs.uk

